20-20-20(all purpose)

Water Soluble Fertilizer

GUARANTEED ANALYSIS

Total Nitrogen (N)	20.00%
Available Phosphate (P ₂ O ₅)	20.00%
Soluble Potash (K ₂ O)	20.00%
\ _ /	0.020%
Boron (B) Copper (Cu) 0.05% Chelated Copper	0.050%
Iron (Fe)	0.100%
0.10% Chelated Iron Manganese (Mn)	0.050%
0.05% Chelated Manganese	0.0005%
Molybdenum (Mo) Zinc (Zn)	0.050%
Chlorine (CI) Maximum	1.000%

Derived From: Ammonium Phosphate, Potassium Nitrate, Urea, Sodium Borate, Copper EDTA, Iron EDTA, Manganese EDTA, Zinc EDTA, and Sodium Molybdate. Chelating Agent is Ethylenediamine Tetra Acetate (EDTA).

DIRECTIONS FOR USE				
	* Rate per 100		Interval Frequency	
Crop	Gal. of Water	Covers	of Application	
Greenhouse Cut Flowers	2 lbs.	100 sq. ft.	10 Days	
Bedding Plants	2 lbs.	100 sq. ft.	7 Days	
Begonias & Tender Plants	1 lb.	100 sq. ft.	10 Days	
Orchards	3 lbs.	Apply to run-off	7-10 Days	

Cucumbers, melons, squash & other vine crops: Use 5 lbs./acre. Begin as soon as the 2nd true leaf unfolds for a total of six (6) applications. **Strawberries**: Use 5 lbs./acre. Begin as soon as growth starts in spring. One (1) to two (2) applications to bloom. Continue after harvest in two (2) to three (3) week intervals.

Potatoes, tomatoes, eggplant, & other solanaceaous plants: Use 5 lbs./acre. In routine crop protection sprays at 7-10 day intervals. **Lawn & Turf:** Use 5 lbs./100 gallons of water for 5000 square feet. Foliar application as growth begins in the spring as needed until dormancy.

CAUTION: THIS PRODUCT CONTAINS BORON, COPPER, IRON, MANGANESE, ZINC, AND MOLYBDENUM. SHOULD BE USED AS RECOMMENDED. MISUSE MAY PROVE HARMFUL. DO NOT USE WITH DINITRO COMPOUNDS, DORMANT OIL, BORDEAUX, SPRAY LIME, OR HIGHLY ALKALINE SPRAY MATERIALS. ALWAYS ADD 20-20-20 TO SPRAY MIXTURE AS THE LAST INGREDIENT.

NET WEIGHT 11.35 KG/25 LBS.

^{*} The rate of 1 lb./100 gallons of water equals 240 ppm Nitrogen.