

Net Wt. 20 lbs. (9.08 kg)

ACTIVE INGREDIENT: EPA Reg. No. 59144-30 EPA Est. 59144-GA-1 (A), 32802-MO-1 (B), 37351-CO-1 (C), 8378-IN-1 (D), 9404-FL-1 (E), 961-PA-01 (F), 961-IL-01 (G), 35497-OR-1 (H), 49991-TX-2 (Q)

IN CASE OF EMERGENCY, CALL 800-424-9300

KEEP OUT OF REACH OF CHILDREN CAUTION See back panel for additional **Precautionary Statements**


PENNINGTON








P.O. Box 290 • Madison, GA 30650



Maximum Application Rates of This Product per Calendar Year By Turf Grass Species $^{(1)}$ This product is recommended for use on the turt grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table: **SETAR LAUNNA MUMIXAM** applications must be made prior to germination of the target weeds. This product will not control established weeds. Base the naximum use rate selection turf species. The length of time of residual weed control provided by this product is related to the rate Apply this product as a single application or in sequential applications to control weeds germinating throughout the year. All

Rates of Application

(6) Do not apply this product to putting greens or areas where dichondra, Colonial bentgrass, velvet bentgrass or annual bluegrass (5) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control. (4) Application to turf stressed by drought, low fertility or pest damage may result in turf injury. (3) Do not cut treated sod within 120 days after application. Do not apply to newly set sod until the following year. (2) Do not apply to overseeded turt within 60 days after seeding or until after the second mowing, whichever is longer. Injury to peciable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus seed in the second inch of soil, not thatch plus soil. (1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses.



USE PRECAUTIONS CRABGRASS SEED GERMINATION DATES



This product is a selective preemergent herbicide that, when properly applied, controls certain grass and broadleaf weeds in established turt grasses and lawns. The maximum amount of product that may be applied per year is given for each turt grass species in the Maximum Annual Rates section of this label. **ESTABLISHED TURF** Uniformly apply using suitable, calibrated application equipment.

Application Directions

Do not graze or feed livestock forage cut from areas treated with this product. Do not apply this product through any type of irrigation system. Do not apply aerially. Do not apply to golf courses or putting greens. Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury or illegal residues. Not for use on plants being grown for sale or other commercial use, commercial seed production or research purposes. turf grasses (excluding golf course putting greens) and lawns. This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control is obtained when this product is activated by at least 1/2 inch of rainfall or irrigation, or shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Directions for Use

This product is a selective preemergent herbicide that provides residual control of many grass and broadleat weeds in established

areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters. This product has low solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal Environmental Hazards

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information. mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. or doctor for treatment advice. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by If In Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on Skin: Take off contaminated clothing, Immediately rinse skin with plenty of water for 15-20 minutes. Call a poison control center If on Skin: Take off contaminated clothing. Immediately rinse skin with plenty of water for 15-20 minutes. Call a poison control center First Aid while handling the material may cause allergic reactions in some individuals.

CAUTION: Causes moderate eye irritation. Harmful if inhaled or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Thoroughly wash with soap and water after handling. Prolonged or frequently repeated skin contact Hazards to Humans and Domestic Animals Precautionary Statements

Derived from: Urea, Ammonium Phosphate, Muriate of Potash. Soluble Potash (K2O). . (₂O₂q) etshqeodq eldslisvA 1.10% Urea Nitrogen Total Nitrogen (N).....3.90% Ammoniacal Nitrogen %00.3 F525 Guaranteed Analysis 5-10-25

COVERS.FT.

+ TYMN EEKLITISEK 2-10-72 CKABGRASS PREVENTER



46" **WEB WIDTH**