



# Anuew<sup>®</sup> EZ

## PLANT GROWTH REGULATOR

**ACTIVE INGREDIENT:**

Prohexadione calcium  
[calcium 3-oxido-5-oxo-4-propionylcyclohex-3-enecarboxylate].....10.0%

**OTHER INGREDIENTS:**.....90.0%

**TOTAL:**.....100.0%

Contains 0.98 lb. Prohexadione calcium per gallon

### KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID, PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE

For Medical Emergencies, Call (877) 325-1840

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

EPA REG. NO. 1001-92

Manufactured by  
Cleary Chemicals, LLC  
11901 S. Austin Ave.  
Alsip, IL 60803

 **Nufarm**  
Grow a better tomorrow



**Net Contents**  
**2.5 Gal.**  
**(9.46 L)**  
**Nonrefillable Container**

15841000

## FIRST AID

<b>IF ON SKIN</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

## HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

#### CAUTION

Harmful if absorbed through the skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

#### Personal Protective Equipment (PPE)

##### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq$  14 mils, nitrile rubber  $\geq$  14 mils, neoprene rubber  $\geq$  14 mils, polyethylene or polyvinyl chloride  $\geq$  14 mils, or viton  $\geq$  14 mils.
- Shoes plus socks

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## USER SAFETY RECOMMENDATIONS

#### User Should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

**DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash waters or rinsate.

## PHYSICAL AND CHEMICAL HAZARDS

**DO NOT** mix or allow contact with oxidizing agents. Hazardous chemical reaction may occur.

#### ENDANGERED SPECIES CONCERNS

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of Federal Law.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq$  14 mils, nitrile rubber  $\geq$  14 mils, neoprene rubber  $\geq$  14 mils, polyethylene or polyvinyl chloride  $\geq$  14 mils, or viton  $\geq$  14 mils.
- Shoes plus socks

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. **DO NOT** enter treated areas without footwear until sprays have dried.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

All applicable directions, restrictions, precautions and **Conditions of Sale and Warranty** are to be followed.

### PRODUCT INFORMATION

ANUEW<sup>®</sup> EZ is a tool for plant growth management. It works by inhibiting late stage biosynthesis of gibberellin, resulting in a decrease in cell elongation, a slowing of vertical growth, and an improvement in plant physiology leading to plant health and stress tolerance.

**Turf:** In turfgrass inhibiting late stage biosynthesis of gibberellin results in a reduction of both the mowing frequency and the amount of grass clippings in the treated area. ANUEW EZ reduces clippings within 2-5 days depending on species. The turfgrass treated with ANUEW EZ grows in a more compact form, often resulting in the enhancement of its color and quality. ANUEW EZ is active through foliar uptake and is not taken up by the plant from the soil therefore, when applying it is important to use enough spray volume to thoroughly wet the turfgrass leaves without significant runoff. A non-ionic spreader to spread spray droplets may be used to assure effective spray coverage. Due to rapid foliar absorption, ANUEW EZ is rainfast within 1-4 hours. ANUEW EZ is most effective when used on actively growing, well-maintained turf. The risk of scalping may be reduced on turfgrass areas treated with ANUEW EZ. It can be applied to golf course fairways, tees, greens and roughs, residential and commercial lawns, sod farms, sports fields, cemeteries, and municipal sites.

**NOTICE TO USER:** Turf tolerance to ANUEW EZ has been found to be acceptable for the turf species listed on this label. Due to the large number of species, cultivars and growing situations it is impossible to test every one for tolerance efficacy and plant safety to ANUEW EZ. Neither the manufacturer nor the seller has determined whether or not ANUEW EZ can be used safely on turf species not specified on this label. For these turf species the professional user may determine if ANUEW EZ can be utilized effectively and safely prior to large scale commercial use. Before using ANUEW EZ on turf species not listed on this label test ANUEW EZ on a small scale first. Start with the lowest directed rate and evaluate the treated turf for plant safety and effective growth regulation prior to widespread use.

## RESTRICTIONS

- **DO NOT** use on food crops.
- **DO NOT** apply ANUEW EZ through any irrigation/chemigation system.
- **DO NOT** apply more than 19.1 gallons (18.7 lb. ai) of ANUEW EZ per acre or 56.1 fl. oz. (0.43 lb.ai) per 1000 square feet per year.
- **DO NOT** apply more than 54 fl. oz. (0.41 lb. ai) of ANUEW EZ per acre or 1.2 fl. oz. (0.009 lb. ai) per 1000 square feet in a single application.
- **DO NOT** make more than 47 applications per year.
- Minimum retreatment: 7 days
- **DO NOT** graze treated areas or feed clippings to livestock.
- **DO NOT** apply ANUEW EZ to grass grown for seed.

## PRECAUTIONS

- Under certain environmental conditions and growth stage conditions on annual bluegrass (*Poa annua*), a transient foliar chlorosis or bronzing may be observed. For optimal results apply to healthy actively growing turfgrass.
- ANUEW EZ may be used on sodded or sprigged turf after it has knitted down and rooted firmly.

## DIRECTIONS FOR USE / APPLICATION METHODS FOR TURFGRASS

ANUEW® EZ may be applied with all types of spray equipment used for ground applications, including properly calibrated boom sprayers, boomless nozzles, spray guns, hand-helds, backpack, mechanical, or motorized spray equipment. Removable chemical extraction probes (also known as “stingers”) used in suction/extraction systems must be rinsed within the pesticide container prior to removal.

Apply ANUEW EZ using 1-4 week intervals based upon turf species, turf growth as influenced by weather conditions, and management to achieve the desired level of growth regulation and playing conditions. See the ANUEW EZ Application Rates Table for product use and application rates, which can be used for growth management and turfgrass color and quality enhancement.

For increased and longer growth suppression, when temporary discoloration can be tolerated, a maximum of twice the ANUEW EZ rate from the Application Rates Table may be applied.

ANUEW EZ performance can be affected by many factors, including environmental conditions, turf vigor and growth stage, soil moisture, fertility levels, and other cultural practices that affect plant growth.

### Application Timing:

Apply ANUEW EZ when the turf is green and actively growing. Delay the application or use a lower rate of ANUEW EZ as designated in the ANUEW EZ Application Rates Table if the turf is entering stressful growing conditions, including pest pressure, high temperatures, low moisture conditions or certain cultural practices including, but not limited to, aeration or verticutting. Repeat applications of ANUEW EZ may be made as noted in the ANUEW EZ Application Rates Table.

Environmental and management conditions can influence turf growth and the potential for uneven regulation. Rapid periods of rebound-growth can be managed by utilizing Growing Degree Day [GDD] modeling based upon climatic conditions at your site. Research has shown that GDD models can be used for re-application timings for common golf situations to assure even turf regulation and to avoid rebound growth.

### Spray Preparation:

Apply ANUEW EZ in a sufficient volume of water to provide uniform and complete wetting of the turfgrass foliage. 0.25 to 5 gallons water/1,000 sq ft may be used, however, for best results 1-2 gallon water/1,000 sq. ft. is advised.

**Adjuvants:** Use of a non-ionic surfactant in the spray mixture may improve coverage of the turf foliage and product performance consistency. Follow the manufacturer's label rate specification.

**Hard Water:** If the water source used for the spray mixture is high (greater than 140 ppm or 140 mg/l) in calcium, add 1 lb. of ammonium sulfate for every pint of ANUEW EZ used in the spray tank. Use a high quality, spray-grade ammonium sulfate product to avoid nozzle plugging.

**Compatibility Test for Tank Mix:** All possible tank mixes have not been evaluated for safety and/or physical compatibility. Tank mix user assumes all risk.

A jar test can be used to evaluate the physical compatibility of components to be mixed in a spray tank with ANUEW EZ before making an application. Use a clear glass quart jar with lid by mixing components in the order shown below under “Mixing Order” in the same relative proportions as it will be used in the spray tank. Cap the jar and invert 10 cycles after adding each component. After all ingredients have been added, let the mixture stand for 15 minutes. Visually, evaluate the suspension for uniformity and stability. The spray mixture must not have particles that precipitate to the bottom of the jar, nor have any oily films or layers visible at the surface, nor contain flakes, balls, or sludge-appearing particles. If the spray mixture doesn't stay uniform in appearance, it is not compatible, and the combination must not be used in a tank mix.

### Mixing Order:

- **Water** - Fill a clean spray tank half full with water and begin continuous agitation.
- **Products in PVA bags** - Place any product contained in water-soluble PVA bags (WVSB) into the mixing tank. Wait until all water-soluble bags have fully dissolved and the product is evenly mixed in the spray tank before adding any further products.
- **ANUEW EZ and water-dispersible products** (WDG; DF), **wettable powders** (WP), and **suspension concentrates** (SC) - Add to the spray tank while maintaining agitation
- **Adjuvants** - Includes non-ionic surfactants
- **Water-soluble products** - Includes micro emulsion products
- **Emulsifiable concentrates** (EC)
- **Water-soluble additives** (S)
- **Remaining quantity of water**

Agitate spray tank mixture constantly during application. Prepare sufficient quantity of product for the immediate application needs. The spray tank mix must not be allowed to sit for prolonged periods without agitation. DO NOT leave product mix in the tank overnight.

Calcium or boron-containing products must not be added to the spray mixture in excess of 140 ppm concentration. Consider hardness of source water as well for these additions. These products will reduce the activity of ANUEW EZ.

**Irrigation after application:** Do not irrigate treated area for 4 hours application.

**Program Scheduling:** Turfgrass areas treated with ANUEW EZ must continue to be maintained normally using proper irrigation, fertility, and pest control product(s) when necessary. Precaution is advised when tank mixing with ANUEW EZ. The spray mixture must be applied on a small area first as a test before treating larger areas.

**Conditions and Practices for Best Performance of ANUEW® EZ:**

- Treated turfgrass is actively growing.
- Use a non-ionic surfactant to help improve leaf coverage and plant uptake of the active ingredient.
- Maximize the time the applied spray mixture remains as a moist film on the turfgrass foliage by avoiding treatments during warm, low humidity, and windy conditions. Longer drying times enhance absorption of ANUEW EZ.

**TABLE 1: ANUEW EZ APPLICATION RATES USING CALENDAR OR GDD MODELING  
TURF USES EXCEPT CALIFORNIA**

	Slows Vertical Growth, Reduces Mowing Intervals, Improves Color and Quality			
	FAIRWAYS AND HIGHER CUT TURF		GREENS AND TEES	
	fl. oz./acre	fl. oz./1,000 sq. ft.	fl. oz./acre	fl. oz./1,000 sq. ft.
<b>Warm Season Turf</b>				
Common and Hybrid Bermudagrass cultivars including 419, Tifway, TifSport, TifEagle, TifGreen, Champion, MiniVerde, and Riviera	18-54 (0.14-0.41 lb. ai)	0.4 – 1.2 (0.003-0.009 lb. ai)	18-36 (0.14-0.28 lb. ai)	0.4 – 0.8 (0.003-0.006 lb. ai)
Zoysiagrass cultivars including Empire, Zenith, Emerald, El Toro and Meyer St. Augustine sp. Paspalum sp. Kikuyugrass sp.	18-54 (0.14-0.41 lb. ai)	0.4 – 1.2 (0.003-0.009 lb. ai)		
<b>Cool Season Turf</b>				
Bentgrass cultivars including: L-93, Penn A4, Penn G2, Pennlinks, Penneagle, Southshore, Declaration, Independence and SR series	9-27 (0.07-0.21 lb. ai)	0.2 – 0.6 (0.002-0.005 lb. ai)	4.5-18 (0.03-0.14 lb. ai)	0.1 – 0.4 (0.0008-0.003 lb. ai)
Annual and Perennial <i>Poa annua</i> cultivars, <i>Poa trivialis</i>	2.2 - 18 (0.02-0.14 lb. ai)	0.05 – 0.4 (0.0004-0.003 lb. ai)	2.2 – 18 (0.02-0.14 lb. ai)	0.05 – 0.4 (0.0004-0.003 lb. ai)
Kentucky bluegrass cultivars including Midnight, Odyssey, Merion	18-36 (0.14-0.28 lb. ai)	0.4 – 0.8 (0.003-0.006 lb. ai)		
Perennial ryegrass cultivars including Manhattan IV Tall Fescues Fine Fescue	18-36 (0.14-0.28 lb. ai)	0.4 – 0.8 (0.003-0.006 lb. ai)		

**TABLE 2: ANUEW® EZ GROWING DEGREE DAY [GDD\*] VALUES  
TURF USES EXCEPT CALIFORNIA**

Common	Scientific	Greens	Fairway
<b>Cool Season Turf</b>		Duration [GDD]	Duration [GDD]
Annual bluegrass	<i>Poa annua</i>	250 - 280	280 - 330
Kentucky bluegrass	<i>Poa pratensis</i>		280 - 310
Bentgrass	<i>Agrostis</i> spp.	260 - 280	350 - 380
Ryegrass	<i>Lolium</i> spp.		280 - 310
<b>Warm Season Turf</b>			
Bermudagrass	<i>Cynodon</i> spp.	125	180 - 225
Zoysiagrass	<i>Zoysia</i> spp.	300	400 - 450
Bahiagrass	<i>Paspalum notatum</i>		180 - 225
Kikuyugrass	<i>Pennisetum clandestinum</i>		290

\*GDDs for cool season turfgrass species are calculated in Celsius with a base of 0°C while GDDs for warm season turfgrass species are calculated in Celsius with a base of 10°C.

**TABLE 3: FOR OVERSEEDING ENHANCEMENT IN BERMUDAGRASS  
TURF USES EXCEPT CALIFORNIA**

Turf Type	fl. oz./acre	Application Interval	Notes
Common and Hybrid Bermudagrass cultivars including 419, Tifway, TifSport, TifEagle, TifGreen, MiniVerde, Riviera, Yukon	2.2 - 4.5 (0.02-0.03 lb. ai)	Apply ANUEW EZ once 3-5 days prior to seeding. Any preparation of the site, including verticutting, spiking, or scalping, must be done 1-2 days after application of ANUEW EZ.	ANUEW EZ is a foliar-active product; therefore, germination and seedling growth of the overseeding are not affected when used as directed. Use good cultural practices in conjunction with this product, including fertilization and irrigation, to optimize results. Some temporary yellowing of the bermudagrass turf may be observed, however, this will not affect the overseeding.

**TABLE 4: FOR GROWTH MANAGEMENT AND COLOR AND QUALITY ENHANCEMENT ON TURF GROWN FOR SOD  
TURF USES EXCEPT CALIFORNIA**

Turf Type	fl. oz./acre	Application Interval	Notes
Warm Season Turf including, but not limited to, Bermudagrass, St. Augustine, etc.	27-54 (0.21-0.41 lb. ai)	2 - 4 week interval	Since turfgrass response can be influenced by a variety of factors, it is advised that the initial evaluation of ANUEW EZ must be limited to a small area. The acceptability of the turfgrass response must be determined using a lower use rate and shorter interval before gradually moving to higher rates and longer intervals.
Cool Season Turf including, but not limited to, Kentucky bluegrass, perennial ryegrass, fescues	18-36 (0.14-0.28 lb. ai)	2 - 4 week interval	

## SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

### Importance of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure specified for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### **Boom Height – Ground Boom**

For ground equipment, the boom must remain level with the crop and have a minimal bounce.

### **Shielded Sprayers**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### **Temperature and Humidity**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### **Temperature Inversions**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

### **Wind**

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

## **STORAGE AND DISPOSAL**

DO NOT contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry place. DO NOT remove the product from the container except for immediate use.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

### **CONTAINER HANDLING:**

**NOTE:** This product is available in multiple containers. Refer to the Net Contents section of this products labeling for the applicable “No refillable” or “Refillable” designation. Follow the container handling instructions below that apply to your container type / size.

**Non-refillable Plastic Containers 5 Gallons or Less:** Non-refillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

**Non-refillable Plastic Containers Larger than 5 Gallons:** Non-refillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable Plastic Containers Larger than 5 Gallons:** Refillable container. Refill this container with pesticide only. DO NOT reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Refillable Plastic Container:** Refill this container with pesticide only. DO NOT reuse this container for any other purpose. Close all openings and replace all caps. Contact Nufarm’s Customer Service Department at 1-800-345-3330 to arrange for return of the empty refillable container.

## **NOTICE OF CONDITIONS OF SALE AND WARRANTY AND LIABILITY LIMITATIONS**

Important: Read the entire Directions for Use and Notice of Conditions of Sale and Warranty and Liability Limitations before using this product. If terms are not acceptable return the unopened container for a full refund.

Our directions for use of this product are based on tests believed to be reliable. However, it is impossible to eliminate all risk associated with the use of this product. Crop injury, inadequate performance, or other unintended consequences may result due to soil or weather conditions, off target movement, presence of other materials, method of use or application, and other factors, all of which are beyond the control of Cleary Chemicals, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer and User.

Cleary Chemicals, LLC warrants that this product conforms to the specifications on the label when used in strict conformance with Direction for Use, subject to the above stated risk limitations. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CLEARY CHEMICALS, LLC MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CLEARY CHEMICALS, LLC'S EXCLUSIVE LIABILITY FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, OR ANY OTHER LEGAL THEORY IS STRICTLY LIMITED TO THE PURCHASE PRICE PAID OR REPLACEMENT OF PRODUCT, AT CLEARY CHEMICALS, LLC'S SOLE DISCRETION.

RV032023

ANUEW is a registered trademark of Nufarm Americas Inc.

All other trademarks are the property of their respective owners.



# Anuew<sup>®</sup> EZ

## PLANT GROWTH REGULATOR

PULL HERE TO OPEN ↑

**ACTIVE INGREDIENT:**

Prohexadione calcium  
[calcium 3-oxido-5-oxo-4-propionylcyclohex-3-enecarboxylate].....10.0%

**OTHER INGREDIENTS:**.....90.0%

**TOTAL:** .....100.0%

Contains 0.98 lb. Prohexadione calcium per gallon

### KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta,  
busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label,  
find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS  
AND DIRECTIONS FOR USE

For Medical Emergencies, Call (877) 325-1840  
For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through the skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

#### FIRST AID

<b>IF ON SKIN</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>

#### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

### STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry place. DO NOT remove the product from the container except for immediate use.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinseate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

#### CONTAINER HANDLING:

**NOTE:** This product is available in multiple containers. Refer to the Net Contents section of this products labeling for the applicable "No refillable" or "Refillable" designation. Follow the container handling instructions below that apply to your container type / size.

**Non-refillable Plastic Containers 5 Gallons or Less:** Non-refillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinseate into application equipment or a mix tank or store rinseate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

**Non-refillable Plastic Containers Larger than 5 Gallons:** Non-refillable container. DO NOT reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinseate into application equipment or a mix tank or store rinseate for later use or disposal. Repeat this procedure two more times. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinseate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable Plastic Containers Larger than 5 Gallons:** Refillable container. Refill this container with pesticide only. DO NOT reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinseate into application equipment or rinseate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

**Refillable Plastic Container:** Refill this container with pesticide only. DO NOT reuse this container for any other purpose. Close all openings and replace all caps. Contact Nufarm's Customer Service Department at 1-800-345-3330 to arrange for return of the empty refillable container.

EPA REG. NO. 1001-92

**Net Contents: 2.5 Gal (9.46 L)**

Manufactured by Cleary Chemicals, LLC  
11901 S. Austin Avenue | Alsip, IL 60803

RV032023[1]A PR