



chipco  
**TRITON<sup>®</sup> 70 WDG**

## Fungicide

*For control of diseases on turfgrasses of commercial, institutional and residential turfgrass, such as golf courses, sod farms, commercial lawns, rights of way, parks, recreation areas, cemeteries and sports fields. Not for use by homeowners.*

ACTIVE INGREDIENT: Triticonazole:.....	70.0%
OTHER INGREDIENTS: .....	30.0%
TOTAL: .....	100.0%

EPA Reg. No. 432-4877 EPA Est. No. indicated by the 3rd and 4th digits of the batch number on this package. (01,03,09,14 or 19)=264-MO-003 (98)=33967-NJ-1 (65)=432-TX-1

**KEEP OUT OF REACH OF CHILDREN  
 CAUTION**

Net Weight  
**2 Pounds**

79400195  
 79365217B 100709AV1

**BACKED  
 by BAYER<sup>™</sup>**

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY**  
Call 24 Hours A Day 1-800-334-7577  
For **PRODUCT USE** Information Call 1-800-331-2867

## FIRST AID

<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Immediately call a poison control center or doctor for treatment advice.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>

**For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577.**  
**Have the product container or label with you when calling a poison control center or doctor or going for treatment.**

**NOTE TO PHYSICIAN:** No specific antidote is available. Treat symptomatically.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

## **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Applicators and other handlers of this product must wear long-sleeved shirt and long pants, chemical-resistant gloves made out of any waterproof material, and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### **User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

## **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water. Drift or runoff may be hazardous to aquatic invertebrates in neighboring areas.

## **DIRECTIONS FOR USE**

**It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**Read entire label before using this product.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

### **AGRICULTURAL USE REQUIREMENTS (continued)**

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls, chemical-resistant gloves made out of any waterproof material, and shoes plus socks.

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

### **STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

#### **PESTICIDE STORAGE**

Store in original container. Keep container closed. Store in an area that is out of reach of children and animals, away from food, feedstuffs, fertilizers and seed. Protect from frost and freezing.

#### **PESTICIDE DISPOSAL**

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### **CONTAINER DISPOSAL**

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Offer for reconditioning, if appropriate. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities.

## GENERAL INFORMATION

Chipco TRITON® 70 WDG Fungicide is a broad spectrum fungicide with preventive, systemic, residual and curative properties recommended for the control or suppression of certain turf diseases. Disease control is optimal when Chipco TRITON 70 WDG Fungicide is applied in a preventive and scheduled spray program and used in rotation with other fungicides. Failure to follow these label instructions may result in turfgrass injury and inferior turfgrass quality.

Chipco TRITON 70 WDG Fungicide can be used on all commercial, institutional and residential turfgrass, such as golf courses, sod farms, commercial and residential lawns, rights of way, parks, recreation areas, cemeteries and sports fields. Not for use by homeowners.

## APPLICATION INFORMATION

Chipco TRITON 70 WDG Fungicide may be used alone to control important diseases of turfgrass or alternatively used in tank mix or sequentially with other registered turfgrass fungicides.

Applications using sufficient water volume to provide thorough and uniform coverage generally provide the most effective disease control. Do not make applications when conditions favor drift beyond the target application area. All applications of Chipco TRITON 70 WDG Fungicide must be made in accordance with the use directions of this label.

Chipco TRITON 70 WDG Fungicide, when used in conjunction with best turfgrass management practice, will control the diseases listed under **TURF-SPECIFIC USE DIRECTIONS**.

Chipco TRITON 70 WDG Fungicide, when combined with **Chipco® Signature™**, will improve disease control and turf quality.

### Turfgrass Species:

Colonial and Creeping Bentgrass  
Common and Hybrid Bermudagrass  
Annual, Rough and Kentucky Bluegrass  
Buffalograss  
Centipedegrass  
Dichondra

Fine and Tall Fescue  
Seashore Paspalum  
Annual and Perennial Ryegrass  
St. Augustinegrass  
Zoysiagrass  
Kikuyugrass

**DO NOT** use on ultradwarf bermudagrass varieties.

**AVOID the shorter application** intervals if the turf is under excessive stress in the summer or under an intensive growth control program.

### Ground Application

For ground application equipment use the appropriate product quantities described in this label in 1 to 5 gallons of water per 1,000 sq ft for turf (44 to 220 gallons per acre).

## **SPRAY MIXING AND COMPATIBILITY**

In general, begin with clean spray equipment and add one-half of the required quantity of water to the spray or mixing tank and start agitation. Refer to the product amounts tabulated under **PRODUCT QUANTITY (OZ) by SPRAY VOLUME and TANK CAPACITY** and add the required quantity of fungicide and the tank-mix partner if applicable to the water and complete filling with water to the required total volume. Follow the recommendations of your State Cooperative Extension Service for tank-mixing with other products. In general, follow the order beginning first with water soluble packaging (wait for it to completely dissolve), wettable powders and water-dispersible granular products, liquid flowables and suspension concentrates and emulsifiable concentrates last. Maintain agitation throughout spraying. Do not allow spray mixture to remain in the tank overnight, or for long periods during the day without agitation.

The turf safety of all potential tank-mixes with Chipco TRITON 70 WDG Fungicide including additives and other pesticides has not been tested on all turf type variants. Before applying any tank-mixture not specifically recommended on this label, safety to the target turf should be confirmed. Chipco TRITON 70 WDG Fungicide is compatible with most commonly used fungicide, herbicide, insecticide, growth regulator and foliar nutrient products. However, the physical compatibility of Chipco TRITON 70 WDG Fungicide with all potential tank-mix partners has not been fully investigated. If tank mixing with other pesticides is desirable, conduct a jar test with the volumes and rates typically used in agricultural application. Using a small container of water, add the proportionate amounts of the products: wettable powders and water-dispersible granular products first, then liquid flowables and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 15 minutes. Look for signs of separation, globules, sludge, flakes, or other precipitates. Physical compatibility is indicated if the combination remains mixed or can be remixed readily.

- The use of a soil penetrating adjuvant may improve the movement into the soil when irrigation after treatment is required for disease control.
- Combinations of Chipco TRITON 70 WDG Fungicide at high labeled rates with PGRs may impact turf quality and reduce turf growth particularly during period of heat stress and high humidity.
- When preparing spray mixtures of Chipco TRITON 70 WDG Fungicide with Bayer Products add the products in the following order: 1) Water dispersible products such as Chipco TRITON 70 WDG Fungicide or Compass® followed by 2) suspension concentrates such as 26 GT®, and lastly 3) Chipco® Signature™.
- This product cannot be mixed with any product whose label contains a prohibition against mixing.

## RESISTANCE MANAGEMENT

The active ingredient in Chipco TRITON 70 WDG Fungicide belongs to the Sterol Inhibitors or Demethylation Inhibitors class of chemistry (Group 3). To maintain long term effectiveness of this fungicide follow the specific resistance management guidance or consult local extension specialist.

To limit the potential for development of disease resistance to this fungicide class do not make more than 3 sequential applications of Chipco TRITON 70 WDG Fungicide or any Group 3 containing fungicide before rotating with a fungicide from a different group.

## GENERAL RESTRICTIONS AND LIMITATIONS

**DO NOT** apply more than the maximum seasonal rate from any combination of products containing triticonazole.

**DO NOT** apply more than 1.8 oz of Chipco TRITON 70 WDG Fungicide per 1,000 sq ft (3.451 lb ai/A) per year.

Follow the **TURF-SPECIFIC USE DIRECTIONS** for repeat application intervals.

For golf courses only: **DO NOT** apply to turf cut higher than 1" on golf holes where water bodies are present.

**This product cannot be used to formulate or reformulate any other pesticide product.**

**DO NOT** apply this product to crops other than turfgrass.

**DO NOT** apply to areas likely to be grazed by livestock.

**DO NOT** feed clippings to livestock or poultry.

**DO NOT** apply through irrigation equipment of any type.

**DO NOT** use on ultradwarf bermudagrass varieties.

## TURF-SPECIFIC USE DIRECTIONS<sup>1</sup>

**CHIPCO TRITON 70 WDG FUNGICIDE controls the diseases listed below.**

Disease Control	Rate oz Product /10,000 sq ft	Use Information
Anthracnose ( <i>Colletotrichum cereale</i> )	0.15 - 0.6	Begin fungicide applications preventively when conditions are favorable for disease development. Continue as needed, on a 14-28 day interval. <b>Under conditions for severe disease or for an early curative application, use the higher rate and shorter interval.</b>

**TURF-SPECIFIC USE DIRECTIONS<sup>1</sup> (continued)**  
**CHIPCO TRITON 70 WDG FUNGICIDE controls the diseases listed below.**

Disease Control	Rate oz Product /1,000 sq ft	Use Information
Brown Patch ( <i>Rhizoctonia solani</i> )	0.225 - 0.6	Begin fungicide applications preventively when conditions are favorable for disease development. Continue as needed, on a 14-28 day interval. <b>Under conditions for severe disease or for an early curative application</b> , use the higher rate and shorter interval.
Brown Patch, Cool weather/Yellow patch ( <i>Rhizoctonia cerealis</i> )	0.3 - 0.6	Make 1 to 2 applications in the fall using a 21-28 day interval or when conditions are favorable for disease development.
Brown Ring (Waitea) Patch ( <i>Waitea circinata</i> var. <i>circinata</i> )	0.3 - 0.6	Begin fungicide applications at the early stage of yellow ring development when conditions are favorable for disease development. Continue as needed, on a 14-28 day interval. Late curative applications will not be effective. <b>Irrigate to move the fungicide beneath the thatch.</b>
Dollar Spot ( <i>Sclerotinia homeocarpa</i> )	0.15 - 0.6	Begin fungicide applications preventively. Continue as needed, on a 14-28 day interval. <b>Under preventive applications</b> where light disease pressure is anticipated, use the lower rate and longer interval. <b>Under conditions for severe disease or for an early curative application</b> , use the higher rate and shorter interval. Rotation with 26 GT® fungicide is suggested for resistance management.



**TURF-SPECIFIC USE DIRECTIONS<sup>1</sup> (continued)****CHIPCO TRITON 70 WDG FUNGICIDE controls the diseases listed below.**

<b>Disease Control</b>	<b>Rate oz Product /1,000 sq ft</b>	<b>Use Information</b>
Fairy Ring	0.3 - 0.6 preventive only	Applications should be made in the spring when 5-day average daily soil temperatures taken at 2 inches are 55 – 65°F. In sandy soils, average air temperature can also be an adequate measure. The second application should be made 28 days later, and applications should be made in 2 gallons H <sub>2</sub> O/1000 ft <sup>2</sup> . In addition, applications should be watered in with 1/8 to 1/4 inch of irrigation. If possible, irrigation should be applied immediately. At the latest, irrigation should occur that night. Chipco TRITON 70 WDG Fungicide applications that are not watered-in have resulted in unsatisfactory control.
Fusarium Patch ( <i>Microdochium nivale</i> )	0.15 - 0.6	Begin fungicide applications preventively in the spring and early summer when the turf stays moist and temperatures are in the range 32-65° F. Repeat applications at 14 day intervals when heavy disease pressure is anticipated. <b>Under conditions for severe disease</b> use the higher rate and tank mix with 26 GT fungicide.

**TURF-SPECIFIC USE DIRECTIONS<sup>1</sup> (continued)**  
**CHIPCO TRITON 70 WDG FUNGICIDE controls the diseases listed below.**

Disease Control	Rate oz Product /1,000 sq ft	Use Information
Gray Snow Mold, Typhula Blight ( <i>Typhula</i> spp.)	0.15 - 0.6	Begin fungicide applications preventively in the late fall. Make 1-2 applications, repeating at 14-28 day intervals when heavy disease pressure is anticipated. <b>Under conditions for severe disease</b> use the higher rate and tank mix with Tartan™ or 26 GT fungicide.
Leaf Spots, including Red Leaf Spot and Gray Leaf Spot ( <i>Dreschlera</i> spp., <i>Bipolaris</i> spp., <i>Helminthosporium</i> spp., <i>Pyricularia</i> spp.)	0.15 - 0.6	Begin fungicide applications preventively. Make 1-2 applications, repeating at 14-28 day intervals when conditions are favorable for development. <b>Under preventive applications</b> where light disease pressure is anticipated, use the lower rate and longer interval. <b>Under conditions for severe disease or for an early curative application</b> , use the higher rate and shorter interval.
Necrotic Ring Spot ( <i>Ophiosphaerella korrae</i> )	0.3 - 0.6	Begin fungicide applications preventively when soil temperature reach 60° F in early spring or summer and continue at 28 day intervals when conditions are favorable for development. <b>Irrigate or aerate to allow movement of fungicide into the root zone.</b>

**TURF-SPECIFIC USE DIRECTIONS<sup>1</sup> (continued)**  
**CHIPCO TRITON 70 WDG FUNGICIDE controls the diseases listed below.**

Disease Control	Rate oz Product /1,000 sq ft	Use Information
Pink Patch ( <i>Limonomyces roseipellis</i> )	0.15 - 0.6	Begin fungicide applications preventively when night air temperatures reach 60° F to 70° F under periods of high rainfall when conditions are favorable for development. <b>Under preventive applications</b> where light disease pressure is anticipated, use the lower rate and longer interval. <b>Under conditions for severe disease or for an early curative application</b> , use the higher rate and shorter interval. Pink Snow Mold
( <i>Microdochium nivale</i> )	0.225 - 0.6	Begin fungicide applications preventively in the late fall just prior to snow cover. Make 2 applications, repeating at a 14-28 day interval when heavy disease pressure is anticipated. <b>Under conditions or prior history for severe disease</b> use the higher rate and tank mix with Tartan or 26 GT fungicide.
Red Thread ( <i>Laetisaria fuciformis</i> )	0.15 - 0.3	Begin fungicide applications preventively when night air temperatures reach 60° F to 70° F under periods of high rainfall when conditions are favorable for development. <b>Under preventive applications</b> where light disease pressure is anticipated, use the lower rate and longer interval. <b>Under conditions for severe disease or for an early curative application</b> , use the higher rate and shorter interval.

**TURF-SPECIFIC USE DIRECTIONS<sup>1</sup> (continued)**  
**CHIPCO TRITON 70 WDG FUNGICIDE controls the diseases listed below.**

Disease Control	Rate oz Product /1,000 sq ft	Use Information
Rust ( <i>Puccinia</i> spp.)	0.15 - 0.3	Begin fungicide applications preventively when conditions are favorable for disease development. Continue as needed, on a 14-28 day interval. <b>Under preventive applications</b> where light disease pressure is anticipated, use the lower rate and longer interval. <b>Under conditions for severe disease or for an early curative application</b> , use the higher rate and shorter interval
Summer Patch ( <i>Magnaporthe poae</i> )	0.3 - 0.6	Begin fungicide applications preventively when soil temperatures reach 60° F to 65° F at a 2-inch soil depth unless otherwise advised by local recommendations. Continue under active disease conditions as needed, on a 14-28 day interval. Use the higher rate and shorter interval under severe disease pressure or if there is prior history.
Take-all Patch ( <i>Gaeumannomyces graminis</i> var. <i>avenae</i> )	0.3 - 0.6	Begin fungicide applications preventively in the fall and repeat in the following spring. Make 1 to 2 applications depending on local disease conditions, repeat under active disease conditions as needed, on a 14-28 day interval. <b>Under conditions for severe disease or for an early curative application</b> , use the higher rate and shorter interval.

**TURF-SPECIFIC USE DIRECTIONS<sup>1</sup> (continued)****CHIPCO TRITON 70 WDG FUNGICIDE controls the diseases listed below.**

<b>Disease Control</b>	<b>Rate oz Product /1,000 sq ft</b>	<b>Use Information</b>
Large Patch ( <i>Rhizoctonia</i> Large Patch Zoysia Patch) ( <i>Rhizoctonia solani</i> )	0.3 - 0.6	Begin fungicide applications preventively in the fall. Make 1-2 applications, repeating at 14-28 day intervals when conditions are favorable for development. <b>Under preventive applications</b> where light disease pressure is anticipated, use the lower rate and longer interval. <b>Under conditions or prior history for severe disease or for an early curative application</b> , use the higher rate and shorter interval.

<b>Additional Turfgrass Uses</b>	<b>Rate oz /1,000 sq ft</b>	<b>Use Information</b>
Algae	0.15 - 0.6	Use preventively the lower rate for the suppression of algae. Continue as needed, on a 14-28 day interval. <b>Tank mix with chlorothalonil or mancozeb to improve the curative control of algae</b>
Summer stress syndrome/summer decline	0.15 - 0.6	Begin applications prior to the onset of conditions for abiotic stress to reduce symptoms of summer stress/decline. Continue as needed, on a 14-28 day interval. <b>Tank mix Chipco Signature to improve disease control and turf quality under moderate to severe abiotic stress.</b>

**<sup>1</sup>DO NOT apply more than 1.8 oz per 1,000 sq ft of CHIPCO TRITON 70 WDG FUNGICIDE per year.**

## PRODUCT QUANTITY (OZ) BY SPRAY VOLUME AND TANK CAPACITY

**CHIPCO TRITON 70 WDG FUNGICIDE at 0.15 ounce per 1000 sq ft.**

**SPRAY VOLUME (gallons per 1000 sq ft)**

	1 Gallon	2 Gallons	3 Gallons	4 Gallons	5 Gallons	
<b>Spray Tank Capacity</b>	<b>25 Gallons</b>	3.8 oz	1.9 oz	1.3 oz	0.94 oz	0.75 oz
	<b>50 Gallons</b>	7.5 oz	3.8 oz	2.5 oz	1.9 oz	1.5 oz
	<b>100 Gallons</b>	15 oz	7.5 oz	5 oz	3.8 oz	3 oz
	<b>200 Gallons</b>	30 oz	15 oz	10 oz	7.5 oz	6 oz

**CHIPCO TRITON 70 WDG FUNGICIDE at 0.23 ounce per 1000 sq ft.**

**SPRAY VOLUME (gallons per 1000 sq ft)**

	1 Gallon	2 Gallons	3 Gallons	4 Gallons	5 Gallons	
<b>Spray Tank Capacity</b>	<b>25 Gallons</b>	5.6 oz	2.8 oz	1.9 oz	1.4 oz	1.1 oz
	<b>50 Gallons</b>	11.3 oz	5.6 oz	3.8 oz	2.8 oz	2.25 oz
	<b>100 Gallons</b>	22.5 oz	11.3 oz	7.5 oz	5.6 oz	4.5 oz
	<b>200 Gallons</b>	45 oz	22.5 oz	15 oz	11.3 oz	9 oz

**CHIPCO TRITON 70 WDG FUNGICIDE at 0.30 ounce per 1000 sq ft.**  
**SPRAY VOLUME (gallons per 1000 sq ft)**

	1 Gallon	2 Gallons	3 Gallons	4 Gallons	5 Gallons
<b>Spray Tank Capacity</b> 25 Gallons	7.5 oz	3.8 oz	2.5 oz	1.9 oz	1.5 oz
50 Gallons	15 oz	7.5 oz	5 oz	3.8 oz	3 oz
100 Gallons	30 oz	15 oz	10 oz	7.5 oz	6 oz
200 Gallons	60 oz	30 oz	20 oz	15 oz	12 oz

**CHIPCO TRITON 70 WDG FUNGICIDE at 0.60 ounce per 1000 sq ft.**  
**SPRAY VOLUME (gallons per 1000 sq ft)**

	1 Gallon	2 Gallons	3 Gallons	4 Gallons	5 Gallons
<b>Spray Tank Capacity</b> 25 Gallons	7.5 oz	3.8 oz	2.5 oz	1.9 oz	1.5 oz
50 Gallons	15 oz	7.5 oz	5 oz	3.8 oz	3 oz
100 Gallons	30 oz	15 oz	10 oz	7.5 oz	6 oz
200 Gallons	60 oz	30 oz	20 oz	15 oz	12 oz

**IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of

Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer (reg'd), the Bayer Cross (reg'd), 26 GT®, Chipco®, Chipco Triton®, Compass®, Signature™, Tartan™ and Backed by Bayer™ are trademarks of Bayer.



Produced for:

**Bayer Environmental Science**

A Division of Bayer CropScience LP

2 T. W. Alexander Drive

Research Triangle Park, NC 27709



chipco  
**TRITON<sup>®</sup> 70 WDG Fungicide**

*For control of diseases on turfgrasses of commercial, institutional and residential turfgrass, such as golf courses, sod farms, commercial lawns, rights of way, parks, recreation areas, cemeteries and sports fields. Not for use by homeowners.*

ACTIVE INGREDIENT: Triticonazole:.....	70.0%
OTHER INGREDIENTS: .....	30.0%
TOTAL: .....	100.0%

EPA Reg. No. 432-4877 EPA Est. No. indicated by the 3rd and 4th digits of the batch number on this package. (01,03,09,14 or 19)=264-MO-003 (98)=33967-NJ-1 (65)=432-TX-1

**KEEP OUT OF REACH OF CHILDREN  
 CAUTION**

For MEDICAL And TRANSPORTATION Emergencies ONLY  
 Call 24 Hours A Day 1-800-334-7577  
 For PRODUCT USE Information Call 1-800-331-2867

Refer to attached leaflet for complete Directions for Use, First Aid, Precautionary Statements and Storage and Disposal Instructions.

Net Weight

**2 Pounds**

79400195

79365217B 100709AV1

**BACKED**  
 by **BAYER<sup>™</sup>**