

3336°WP

FUNGICIDE

Prevent and Cure.

3336® WP is a broad-spectrum fungicide with preventive and curative properties. Its systemic mode of action is effective on a wide range of troublesome diseases in ornamental settings.

FLEXIBLE APPLICATION OPTIONS

- · Premeasured, water-soluble packaging
- Broad-spectrum disease control
- · Full chemigation label
- Tank-mix compatible with most commonly used pesticides

EPA REG. NO.	1001-63
ACTIVE INGR.	thiophanate-methyl (50%)
FORMULATION	wettable powder
CHEM. FAMILY	thiophanates
FRAC NO.	1
REI	12 hours
PACKAGE SIZE	6 x 2 lb
APPLICATION	spray, drench, dip





GREENHOUSE

CAUTION







FUNGICIDE

KEY USES

Annual and perennial flowers
Bedding plants
Foliage plants
Groundcovers
Deciduous and evergreen trees and shrubs

KEY DISEASES CONTROLLED

(See product label for complete list)

Anthracnose
Botrytis
Brown rot and blight
Cercospora
Ovulinia blight
Powdery mildew
Rhizoctonia
Thielaviopsis

APPLICATION RECOMMENDATIONS

3336® WP is a broad-spectrum systemic fungicide which controls a variety of foliar, stem, and root diseases on a wide range of commercially important plants. 3336 WP is also effective as a pre-plant dip on cuttings and bulbs. For soil drench applications, best crop protection is achieved with preventive treatments repeated every 21-28 days. For foliar applications, begin treatments when disease first appears, or during suspected periods of disease incidence. Apply additional applications every 7-14 days or as otherwise instructed for prevention or control. Use of a wetting agent is recommended for plants that have leaves that are difficult to wet properly. Use of a spreader-sticker is recommended to enhance product performance in wet weather conditions or during periods of overhead irrigation. 3336 WP may be applied as a ground application using hand held, mechanical or motorized spray equipment, or as a chemigation spray or through an applicable sprinkler irrigation system; or as an overhead application where applicable. For foliar applications other than cut flowers do not exceed 3.0 pounds active ingredient per acre per single application. For cut flowers do not exceed 0.5 pound active ingredient per single application. Do not exceed 36 pounds active ingredient per acre per crop season from all thiophanatemethyl containing products. For soil drench applications, do not exceed 300 pounds active ingredient per acre per crop season from all thiophanate-methyl containing products.

APPLICATION INSTRUCTIONS

Apply material with properly calibrated, hand held, mechanical or motorized spray equipment or by chemigation through appropriate sprinkler irrigation, flood, or drip systems. Begin applications when disease first appears and repeat at 7-14 day intervals or as needed during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix 8-24 oz of 3336 WP per 100 gal water and apply as a full coverage spray to drip for prevention and control. Spray volume may range up to 400 gallons of finished spray per acre depending upon plant species and plant growth stage. For applications through irrigation systems, refer to use rates indicated in application chart found on the label.

Do not apply to home orchards/fruit trees after fruit set.

Do not use fruits, nuts or sap from treated trees as food or feed.

See label for complete application rates and recommendations.

