



Label

White papers

Coupons

Other

Image

MERCK ANIMAL HEALTH Intervet Inc.

2 GIRALDA FARMS, MADISON, NJ, 07940

Customer Service: 800-521-5767

Order Desk: 800-648-2118

Technical Service (Companion Animal): 800-224-5318

Technical Service (Livestock): 800-211-3573

Fax: 973-937-5557

Website: www.merck-animal-health-usa.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

SAFE-GUARD® Intervet/Merck Animal Health (fenbendazole)

Medicated Dewormer

for Beef & Dairy Cattle

0.5% (fenbendazole) Alfalfa-Based Pellets

ACTIVE DRUG INGREDIENT

fenbendazole	0.5% (2.27 g/lb)
--------------	------------------

GUARANTEED ANALYSIS

Crude Protein	(min.) 15%
Crude Fat	(min.) 1%
Crude Fiber	(max.) 30%
Moisture Content	(max.) 13%
Ash Content	(max.) 15%

INERT INGREDIENTS

Dehydrated Alfalfa Meal, Roughage Products, Calcium Carbonate, Mineral Oil.

CONSULT YOUR VETERINARIAN FOR ASSISTANCE IN THE DIAGNOSIS, TREATMENT, AND CONTROL OF PARASITISM.

CATTLE: Dairy and beef cattle

For the removal and control of: **Lungworms:** (*Dictyocaulus viviparus*). **Stomach worms:** Barberpole worms (*Haemonchus contortus*), brown stomach worms (*Ostertagia ostertagi*), small stomach worms (*Trichostrongylus axei*). **Intestinal worms:** Hookworms (*Bunostomum phlebotomum*), thread-necked intestinal worms (*Nematodirus helvetianus*), small intestinal worms (*Cooperia punctata* & *C. oncophora*), Bankrupt worms (*Trichostrongylus colubriformis*), Nodular worms (*Oesophagostomum radiatum*).

DAIRY AND BEEF CATTLE DOSAGE

5 mg fenbendazole per kg body weight in a **one (1) day treatment** (2.27 mg fenbendazole per pound of body weight).

Examples of Feeding Rates for Safe-Guard® 0.5% (fenbendazole) Alfalfa-Based Pellets for Cattle

Body Weight (lbs)	Safe-Guard® 0.5% Pellets
200	0.2 lbs
500	0.5 lbs
1000	1.0 lbs

Feed for **one (1) day**. No prior withdrawal of feed or water necessary. When feed containing Safe-Guard® has been fed for 1 day according to the above rates, a total intake of 2.27 mg fenbendazole per pound of body weight is assured.

Under conditions of continued exposure to parasites, retreatment may be needed after 4-6 weeks.

Residue Warning: Cattle must not be slaughtered within 13 days following last treatment. For dairy cattle, there is no milk withdrawal period.

WARNING: KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN. NOT FOR USE IN HUMANS.

Distributed by: **Intervet Inc.**, Millsboro, DE 19966

NET WEIGHT	
10 LBS (4.54 KG)	580215-A

CPN: 1047470.3

By using this site, you agree to the [Terms of Service](#) and [Privacy Policy](#).

Copyright 2020 - Animalytix LLC