

SensiPro™ RPM BLUE

DIRECTIONS FOR USE: Always wear approved eye protection and protective gloves.

SUGGESTED USE RATE FOR SENSIPRO™ BLUE GRANULES AS A SPRAY PATTERN INDICATOR:
1-2 ounces by weight per acre (50 gallons water) or 60-120 grams by weight per hectare (450 liters water). Backpack/Knapsack rate for spot spraying is 5 grams or 1 level teaspoon per 10 liters of water. Determine size of area being treated. Fill spray tank 1/3 full with water and remove strainer basket. With continual tank agitation, slowly add predetermined SensiPro™ Blue Granules. Allow to disperse, and then add other actives and water as needed. Spot test and adjust colorant prior to full scale spraying. Taller grass may require higher rates. Retain usage rate records for future reference.

SUGGESTED USE RATE FOR SENSIPRO™ BLUE GRANULES AS A POND/LAKE COLORANT:
Usage rates begin at 1 ounce per 1/16th acre (4-6 feet deep) or 30 grams per .025 hectare (1-2 meters deep). For use only in ponds/lakes in control of the user.

PRECAUTIONARY STATEMENT: Avoid contact with skin, eyes or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. If swallowed: Call a poison control center or doctor immediately for treatment advice.

STORAGE & DISPOSAL: Store in original container in a cool, dry place. Keep from freezing. Triple rinse, then offer for recycling or dispose of in a sanitary landfill or by other procedures approved by state & local authorities.

057261536
05726153B
1043748

READ & UNDERSTAND LABELS PRIOR TO USE

Operator restricts powered travel and hydraulic function operation when operator leaves normal operating position. Lift trucks with OPSS operate differently than similar lift trucks without OPSS.

OR
ACTION AND

TOYOTA
MODEL 8FB
TYPE
MAY